

Results from the OPERA experiment

DONATO DI FERDINANDO ON BEHALF OF OPERA COLLABORATION

INFN sez. di Bologna

The OPERA long baseline experiment located in the INFN Gran Sasso underground Laboratory was designed to establish the $\nu_\mu \rightarrow \nu_\tau$ oscillations in appearance mode, using the CNGS neutrino beam. Five ν_τ candidate events have been detected in a data sample from the 2008-2012 runs, with a expected background of 0.25 events. The background only hypothesis is rejected with a significance larger than 5σ . The analysis of the tau neutrino sample in the framework of the 3+1 neutrino model is presented. OPERA is able to identify ν_e CC events due to the good tracking capabilities of its target. The electron neutrino sample is also used to set limits on the 3+1 oscillation parameters.

PRESENTED AT

NuPhys2015, Prospects in Neutrino Physics
Barbican Centre, London, UK, December 16–18, 2015

1 The OPERA detector

The OPERA (Oscillation Project with Emulsion tRacking Apparatus) [1] experiment was designed to conclusively prove the existence of $\nu_\mu \rightarrow \nu_\tau$ oscillations by the appearance of ν_τ in a pure ν_μ beam. The detector was located in the underground Gran Sasso Laboratory, 730 km from CERN, on the CNGS neutrino beam line. The ν_τ signature is the detection of a τ -lepton (flight length of about 600 μm) produced by Charged Current (CC) interaction with the target mass of the detector. To cope with the challenging task of keeping a high granularity in a large detector mass the technique of Emulsion Cloud Chamber (ECC) has been used. The OPERA ECC target unit (called *brick*) is composed of a sequence of 57 nuclear emulsion sheets interleaved with 56, 1 mm thick, lead plates for a mass of 8.3 kg; its transverse size is $12.5 \times 10.2 \text{ cm}^2$ and the thickness along the beam direction is about 10 radiation lengths. A nuclear emulsion film is made of a 205 μm plastic base with a 44 μm sensitive layer on both sides. In the OPERA apparatus 150000 bricks were arranged in two target sections having each 28 vertical planes (called *walls*) transverse to the beam; each wall was followed by two planes of plastic scintillator strips that composed the Target Tracker (TT) allowing to locate the brick in which the neutrino interaction occurred with a precision of $\mathcal{O}(\text{cm})$. Each target section was completed by a magnetic spectrometer instrumented with Resistive Plate Chamber (RPC) detectors and high precision Drift Tubes (DT).

2 Event reconstruction and analysis

The electronic detector data was used to identify the brick in which the neutrino interaction occurred. Events were classified as Charged Current like (1μ) if a track was tagged as a muon or if a minimum number of detector walls was traversed, or as Neutral Current like (0μ) in the complementary case [2]. The bricks tagged as containing a neutrino interaction vertex were extracted from the detector, their emulsion films were then developed and sent to the scanning laboratories in Europe and Japan, where a search for τ decays was performed using custom automated optical microscopes. The search for the neutrino interaction vertex starts from a set of track predictions, provided by the electronic detectors, found tracks are followed back in the brick, film by film, from the most downstream one to the point where they originate. When a track disappears (i.e. is not found in three consecutive films) a volume of 1 cm^2 by 5 films upstream and 10 films downstream of the possible vertex point is scanned in order to fully explore the vertex area and possibly find other associated particle tracks. The search for the possible decay of a secondary track is then performed through a dedicated decay search procedure [4]. Vertexes for which one track has an impact parameter larger than 10 μm are further analysed

[2, 3], searching for secondary decay vertexes. According to the decay topology of the event, either in one prong (electron, muon or hadron) or in three prongs, kinematical selection criteria are then applied to validate the candidate for appearance of the τ lepton.

3 $\nu_\mu \rightarrow \nu_\tau$ oscillations search

During the CNGS runs from 2008 to 2012, 19505 neutrino interactions were recorded in the target fiducial volume, corresponding to $17.97 \cdot 10^{19}$ protons on target (p.o.t.). The decay search procedure has been applied, after analysis of most of the data, to 5408 events. Out of this sample, five τ candidates match the reconstruction criteria introduced in the previous section. One of them is in the $\tau \rightarrow \mu$ decay channel [7], three in $\tau \rightarrow 1h$ [5, 8, 9] and one in $\tau \rightarrow 3h$ [6]. In the analysed sample 0.25 ± 0.05 background events are expected, mainly from charmed events with an undetected primary muon, the remaining part coming from hadronic re-interactions (for the hadronic decay channels) and large angle muon scattering (for the $\tau \rightarrow \mu$ channel) as shown in Tab. 1. Taking into account the different signal-to-noise ratio for each decay channel, by the observation of five candidates a 5.1σ significance for the exclusion of the background-only hypothesis in the search for $\nu_\mu \rightarrow \nu_\tau$ oscillation is achieved [9].

Channel	Expected background				Expected signal	Observed
	Charm	Had. re-interac.	Large μ -scat.	Total		
$\tau \rightarrow 1h$	0.017 ± 0.003	0.022 ± 0.006	—	0.04 ± 0.01	0.52 ± 0.10	3
$\tau \rightarrow 3h$	0.17 ± 0.03	0.003 ± 0.001	—	0.17 ± 0.03	0.73 ± 0.14	1
$\tau \rightarrow \mu$	0.004 ± 0.001	—	0.0002 ± 0.0001	0.004 ± 0.001	0.61 ± 0.12	1
$\tau \rightarrow e$	0.03 ± 0.01	—	—	0.03 ± 0.01	0.78 ± 0.16	0
Total	0.22 ± 0.04	0.02 ± 0.01	0.0002 ± 0.0001	0.25 ± 0.05	2.64 ± 0.53	5

Table 1: Expected signal and background events for the analysed data sample.

4 Search for sterile neutrinos

The neutrino oscillation phenomenon is currently well described by the mixing of three neutrino flavor eigenstates (ν_e , ν_μ and ν_τ) with three mass eigenstates (ν_1 , ν_2 and ν_3). Nevertheless there are experimental results that cannot be accomodated within this framework [10]. These anomalies may hint to the existence of sterile neutrino(s) with $\Delta m^2 \sim 1 \text{ eV}^2$. The OPERA experiment can test the sterile neutrino hypothesis looking for deviations from predictions of the standard three-neutrino model. OPERA ν_τ appearance results have been used to derive limits on the mixing parameters of a massive ($\sim \text{eV}$) sterile neutrino [12].

In presence of a fourth sterile neutrino with mass m_4 , the oscillation probability is

a function of the 4×4 mixing matrix U and of the three squared mass differences. Observed neutrino oscillation anomalies, if interpreted in terms of one additional sterile neutrino, suggest $|\Delta m_{41}^2|$ values at the eV^2 scale. In the framework of the 3+1 model, at high values of Δm_{41}^2 , the measured 90% C.L. upper limit on the mixing term $\sin^2 2\theta_{\mu\tau} = 4|U_{\mu 4}|^2|U_{\tau 4}|^2$ is 0.116, independently of the mass hierarchy of the three standard neutrinos. The OPERA experiment extends the exclusion limits on Δm_{41}^2 in the $\nu_\mu \rightarrow \nu_\tau$ appearance channel down to values of 10^{-2}eV^2 at large mixing for $\sin^2 2\theta_{\mu\tau} \gtrsim 0.5$ as shown in Figure 1.




Figure 1: OPERA 90% C.L. exclusion limits in the Δm_{41}^2 vs $\sin^2 2\theta_{\mu\tau}$ parameter space for the normal (NH, dashed red) and inverted (IH, solid blue) hierarchy of the three standard neutrino masses. The exclusion plots by other experiments are also shown. Bands are drawn to indicate the excluded regions.

A systematic search for ν_e events was performed in the 2008-2009 data sample. Out of 505 0μ events with located interaction vertex, corresponding to an integrated intensity of $5.25 \cdot 10^{19}$ p.o.t. [11], 19 ν_e candidate events were observed. This number is compatible with the expected ν_e from the beam contamination (19.8 ± 2.8). The current result on the search for the three-flavour neutrino oscillation yields an upper limit $\sin^2 2\theta_{13} < 0.44$ (90% C.L.).

OPERA limits the parameter space available for a non-standard ν_e appearance suggested by the results of the LSND and MiniBooNE experiments [10]. It further constrains the still allowed region around $\Delta m_{new}^2 = 5 \cdot 10^{-2} \text{ eV}^2$. For large Δm_{new}^2 values, a Bayesian approach has been used and the upper limit on $\sin^2 2\theta_{new}$ reaches the value $7.2 \cdot 10^{-3}$.

These results will be updated using 50 ν_e candidates observed in the full data set.

5 Conclusions

The OPERA experiment taking data from 2008 to 2012, collected $17.97 \cdot 10^{19}$ p.o.t. Five ν_τ candidates have been observed and the observation of $\nu_\mu \rightarrow \nu_\tau$ oscillations due to background only is excluded at 5.1σ .

The observed number of ν_e interactions is compatible with the non-oscillation hypothesis, but allows OPERA to set an upper limit in the parameter space for a non-standard ν_e appearance.

Limits on the mixing parameters of a massive sterile neutrino have also been derived in the $\nu_\mu \rightarrow \nu_\tau$ appearance channel and the exclusion limits on Δm_{41}^2 have been extended down to values of 10^{-2} eV^2 at large mixing for $\sin^2 2\theta_{\mu\tau} \gtrsim 0.5$.

References

- [1] R. Acquafredda et al. (OPERA Collab.), JINST **4** (2009) P04018.
- [2] N. Agafanova et al. (OPERA Collab.), New J. Phys. **13** (2011) 053051.
- [3] N. Agafanova et al. (OPERA Collab.), New J. Phys. **14** (2012) 013026.
- [4] N. Agafanova et al. (OPERA Collab.), Eur. Phys. J C **89** (2014) 2986.
- [5] N. Agafanova et al. (OPERA Collab.), Phys. Lett. B **691** (2010) 138.
- [6] N. Agafanova et al. (OPERA Collab.), JHEP **11** (2013) 036.
- [7] N. Agafanova et al. (OPERA Collab.), Phys. Rev. D **89** (2014) 051102.
- [8] N. Agafanova et al. (OPERA Collab.), Prog. Theor. Exp. Phys. (2014) 101C01.
- [9] N. Agafanova et al. (OPERA Collab.), Phys. Rev. Lett. **115** (2015) 12 121802.
- [10] A. Aguilar-Arevalo et al. [LSND Collaboration], Phys. Rev. D **64** (2001) 112007; A. A. Aguilar-Arevalo et al. [MiniBooNE Collaboration], Phys. Rev. Lett. **110** (2013) 161801; G. Mention et al., Phys. Rev. D **83** (2011) 073006; Th. A. Muller et al., Phys. Rev. C **83** (2011) 054615; C. Giunti and M. Laveder, Phys. Rev. C **83** (2011) 065504.
- [11] N. Agafanova et al. (OPERA Collab.) JHEP **07**(2013) 004.
- [12] N. Agafonova et al. (OPERA Collab.) JHEP **1506** (2015) 069.